

ON EXPONENTIAL STABILITY OF WONHAM FILTER

P. CHIGANSKY AND R. LIPTSER

ABSTRACT. We give elementary proof of a stability result from Atar and Zeitouni, [1], concerning an exponential asymptotic ($t \rightarrow \infty$) for filtering estimates generated by wrongly initialized Wonham filter. This proof is based on new exponential bound having independent interest.

1. INTRODUCTION

In [1], Atar and Zeitouni established an exponential asymptotic (in $t \rightarrow \infty$) rate of convergence for the distance between differently initialized filter outputs when a signal is Markov process valued in a finite state space for both the discrete and continuous time cases (see related topic in [2]). In this note, we discuss only continuous time case¹ and deal with exponential stability of the Wonham filter. Following [1], we assume that a signal $X = (X_t)_{t \geq 0}$ is homogeneous Markov process, valued in a finite alphabet $\mathbb{S} = \{a_1, \dots, a_d\}$, with transition intensities matrix Λ (with entries λ_{ij}) and initial distribution vector ν . An observation $Y = (Y_t)_{t \geq 0}$ is defined as

$$Y_t = \int_0^t h(X_s) ds + \sigma W_t$$

where $W = (W_t)_{t \geq 0}$ is a Wiener process independent of X , σ is a constant and h is $\mathbb{S} \xrightarrow{h} \mathbb{R}$ function. The Wonham filter, [5] (see also Ch. 9 in [4]), creates π_t the conditional distribution of X_t given $Y_{[0,t]} = \sigma\{Y_s, s \leq t\}$, that is $\pi_t(1) = P(X_t = a_1 | Y_{[0,t]})$, $\dots$, $\pi_t(d) = P(X_t = a_d | Y_{[0,t]})$:

$$\begin{aligned} \pi_0 &= \nu \\ d\pi_t &= \Lambda^* \pi_t dt + (\text{diag}(\pi_t) - \pi_t \pi_t^*) h \sigma^{-2} (dY_t - h^* \pi_t dt), \end{aligned} \quad (1.1)$$

where * , $\text{diag}(\cdot)$, and h are the transposition symbol, scalar matrix, and vector with entries $h(a_1), \dots, h(a_d)$ respectively.

It is assumed in [1] that

$$\lambda_{ij} > 0, \quad \forall i \neq j \quad (1.2)$$

and therefore the transition probabilities of Markov process X converge exponentially fast to the unique and positive invariant distribution. In [1], it

1991 *Mathematics Subject Classification.* 93E11, 60J57.

Key words and phrases. Hidden Markov Models, Wonham filter, Exponential stability.

¹regarding to discrete time case we recommend also the recent paper of Le Gland and Mevel [3].

is shown that this property is inherited, in some sense, by π_t . To emphasize the dependence of π_t on ν , below we write π_t^ν instead of π_t . Assume β the probability distribution on $\mathbb{S}$, different from ν , is used in (1.1) as the initial condition and denote $\pi_t^{\beta\nu}$ the solution of corresponding Itô equation from (1.1). For wrong initial distributions β', β'' , denote $\|\pi_t^{\beta'\nu} - \pi_t^{\beta''\nu}\|$ the total variation norm for difference of wrong conditional distributions. The following result is known from [1] (see Theorem 1.6). For any β', β''

$$\overline{\lim}_{t \rightarrow \infty} \frac{1}{t} \log \|\pi_t^{\beta'\nu} - \pi_t^{\beta''\nu}\| \leq -2 \min_{p \neq q} \sqrt{\lambda_{pq} \lambda_{qp}}, \quad P - \text{a.s.} \quad (1.3)$$

This result is very important in applications since it guarantees that wrongly initialized Wonham filter generates a wrong conditional distribution converging to correct one exponentially fast as $t \rightarrow \infty$.

The proof of (1.3) in [1] (see pp. 46-48 there), being the pioneer, uses “heavy” tools such as Oseledec’s multiplicative ergodic theorem, Birghoff’s contraction coefficient, etc. So, it is surprising that an elementary proof of (1.3) is possible and, moreover, an exponential estimate for $\|\pi_t^{\beta\nu} - \pi_t^\nu\|$, with the same rate $-2 \min_{p \neq q} \sqrt{\lambda_{pq} \lambda_{qp}}$, is valid for any $t > 0$.

Theorem 1.1. *Assume $\lambda_{ij} > 0$ for any $i \neq j$. Assume ν is positive and $\beta \sim \nu$. Then, P -a.s. for any $t \geq 0$*

$$\|\pi_t^{\beta\nu} - \pi_t^\nu\| \leq d^2 \max_{j \in \mathbb{S}} \frac{d\beta}{d\nu}(j) \max_{j \in \mathbb{S}} \frac{d\nu}{d\beta}(j) \exp \left(-2 \min_{p \neq q} \sqrt{\lambda_{pq} \lambda_{qp}} t \right). \quad (1.4)$$

2. PROOF OF THEOREM 1.1

For a notational convenience, write X_t^α, Y_t^α (instead of X_t, Y_t), if α is the distribution of X_0 . Then, under ν being the distribution of X_0 , write

$$\pi_t^\nu(i) = P(X_t^\nu = a_i | Y_{[0,t]}^\nu), \quad i = 1, \dots, d.$$

Due to (1.1), $\pi_t^{\beta\nu}$ is defined as

$$\begin{aligned} \pi_0^{\beta\nu} &= \beta \\ d\pi_t^{\beta\nu} &= \Lambda^* \pi_t^{\beta\nu} dt + (\text{diag}(\pi_t^{\beta\nu}) - \pi_t^{\beta\nu} (\pi_t^{\beta\nu})^*) h \sigma^{-2} (dY_t^\nu - h^* \pi_t^{\beta\nu} dt). \end{aligned} \quad (2.1)$$

It is obvious that $\pi_t^{\beta\nu}(i) = P(X_t^\beta = a_i | Y_{[0,t]}^\beta) \Big|_{Y^\beta := Y^\nu}$, $i = 1, \dots, d$.

Set

$$\rho_{ji}(t) = P(X_0^\nu = a_j | Y_{[0,t]}^\nu, X_t^\nu = a_i), \quad i, j = 1, \dots, d. \quad (2.2)$$

2.1. Auxiliary lemmas.

Lemma 2.1. *Under the assumptions of Theorem 1.1, for any $i = 1, \dots, d$*

$$|\pi_t^\nu(i) - \pi_t^{\beta\nu}(i)| \leq d \max_{j \in \mathbb{S}} \frac{d\beta}{d\nu}(j) \max_{j \in \mathbb{S}} \frac{d\nu}{d\beta}(j) \max_{1 \leq i, j, k \leq d} |\rho_{ji}(t) - \rho_{jk}(t)|, \quad t \geq 0.$$

Proof. Denote Q^ν and Q^β distributions of (X^ν, Y^ν) and (X^β, Y^β) respectively. Both (X^ν, Y^ν) and (X^β, Y^β) are Markov processes, with paths from the Skorokhod space $\mathbb{D}_{[0,\infty)}(\mathbb{R}^2)$. They have the same transition probabilities, $Y_0^\nu \equiv Y_0^\beta (= 0)$ and different initial distributions for X_0^ν and X_0^β : ν and β respectively. Since $\nu \sim \beta$, the measures Q^ν and Q^β are equivalent and

$$\frac{dQ^\beta}{dQ^\nu}(X^\nu, Y^\nu) \equiv \frac{d\beta}{d\nu}(X_0^\nu).$$

We show first that for any $i = 1, \dots, d$ and $t > 0$, Q^ν - and Q^β -a.s.

$$\pi_t^{\beta\nu}(i) = \frac{\sum_{j=1}^d \left(\frac{d\beta}{d\nu}(j) P(X_0^\nu = a_j, X_t^\nu = a_i | Y_{[0,t]}^\nu) \right)}{E\left(\frac{d\beta}{d\nu}(X_0^\nu) | Y_{[0,t]}^\nu\right)}. \quad (2.3)$$

Let $(\mathcal{D}_t)_{t \geq 0}$ be a right continuous filtration on $\mathbb{D}_{[0,\infty)}(\mathbb{R}^2)$ completed by sets of Q^ν -measure zero. For fixed $t > 0$, let $\psi(x, y) \equiv \psi(y)$, $(x, y) \in \mathbb{D}_{[0,\infty)}(\mathbb{R}^2)$, be $\mathcal{D}_t$ measurable and bounded function. Then

$$\begin{aligned} E\psi(Y^\nu) \pi_t^{\beta\nu}(i) E\left(\frac{d\beta}{d\nu}(X_0^\nu) | Y_{[0,t]}^\nu\right) &= E\psi(Y^\nu) \pi_t^{\beta\nu}(i) \frac{d\beta}{d\nu}(X_0^\nu) \\ &= E\psi(Y^\nu) \pi_t^{\beta\nu}(i) \frac{dQ^\beta}{dQ^\nu}(X^\nu, Y^\nu) = E\psi(Y^\beta) \pi_t^\beta(i) = E\psi(Y^\beta) I(X_t^\beta = a_i) \\ &= E\psi(Y^\nu) I(X_t^\nu = a_i) \frac{dQ^\beta}{dQ^\nu}(X^\nu, Y^\nu) = E\psi(Y^\nu) I(X_t^\nu = a_i) \frac{d\beta}{d\nu}(X_0^\nu) \\ &= E\psi(Y^\nu) E\left(I(X_t^\nu = a_i) \frac{d\beta}{d\nu}(X_0^\nu) | Y_{[0,t]}^\nu\right). \end{aligned}$$

So, by an arbitrariness of $\psi(Y^\nu)$, we get $\pi_t^{\beta\nu}(i) = \frac{E\left(I(X_t^\nu = a_i) \frac{d\beta}{d\nu}(X_0^\nu) | Y_{[0,t]}^\nu\right)}{E\left(\frac{d\beta}{d\nu}(X_0^\nu) | Y_{[0,t]}^\nu\right)}$

and it remains to notice that

$$E\left(I(X_t^\nu = a_i) \frac{d\beta}{d\nu}(X_0^\nu) | Y_{[0,t]}^\nu\right) = \sum_{j=1}^d \frac{d\beta}{d\nu}(j) P(X_t^\nu = a_i, X_0^\nu = a_j | Y_{[0,t]}^\nu).$$

Now, taking into the consideration (2.3), write

$$\begin{aligned} |\pi_t^\nu(i) - \pi_t^{\beta\nu}(i)| &= \left| \pi_t^\nu(i) - \frac{\sum_{j=1}^d \left(\frac{d\beta}{d\nu}(j) P(X_0^\nu = a_j, X_t^\nu = a_i | Y_{[0,t]}^\nu) \right)}{E\left(\frac{d\beta}{d\nu}(X_0^\nu) | Y_{[0,t]}^\nu\right)} \right| \\ &= \frac{\left| \sum_{j=1}^d \frac{d\beta}{d\nu}(j) \left(\pi_t^\nu(i) P(X_0^\nu = a_j | Y_{[0,t]}^\nu) - P(X_0^\nu = a_j, X_t^\nu = a_i | Y_{[0,t]}^\nu) \right) \right|}{E\left(\frac{d\beta}{d\nu}(X_0^\nu) | Y_{[0,t]}^\nu\right)}. \end{aligned}$$

Notice that by the Jensen inequality $1/E\left(\frac{d\beta}{d\nu}(X_0^\nu)|Y_{[0,t]}^\nu\right) \leq E\left(\frac{d\nu}{d\beta}(X_0^\nu)|Y_{[0,t]}^\nu\right)$. Consequently,

$$\begin{aligned}
|\pi_t^\nu(i) - \pi_t^{\beta\nu}(i)| &\leq \max_{j \in \mathbb{S}} \frac{d\beta}{d\nu}(j) \max_{j \in \mathbb{S}} \frac{d\nu}{d\beta}(j) \\
&\times \left| \sum_{j=1}^d \pi_t^\nu(i) \left(P(X_0^\nu = a_j | Y_{[0,t]}^\nu) - P(X_0^\nu = a_j, X_t^\nu = a_i, Y_{[0,t]}^\nu) \right) \right| \\
&\leq \max_{j \in \mathbb{S}} \frac{d\beta}{d\nu}(j) \max_{j \in \mathbb{S}} \frac{d\nu}{d\beta}(j) \\
&\times \sum_{j=1}^d \pi_t^\nu(i) \left| P(X_0^\nu = a_j | Y_{[0,t]}^\nu) - P(X_0^\nu = a_j | X_t^\nu = a_i, Y_{[0,t]}^\nu) \right| \\
&\leq \max_{j \in \mathbb{S}} \frac{d\beta}{d\nu}(j) \max_{j \in \mathbb{S}} \frac{d\nu}{d\beta}(j) \sum_{j=1}^d \left| P(X_0^\nu = a_j | Y_{[0,t]}^\nu) - P(X_0^\nu = a_j | X_t^\nu = a_i, Y_{[0,t]}^\nu) \right| \\
&= \max_{j \in \mathbb{S}} \frac{d\beta}{d\nu}(j) \max_{j \in \mathbb{S}} \frac{d\nu}{d\beta}(j) \sum_{j=1}^d \left| P(X_0^\nu = a_j | Y_{[0,t]}^\nu) - \rho_{ji}(t) \right|. \quad (2.4)
\end{aligned}$$

Further, an obvious formula $P(X_0^\nu = j | Y_{[0,t]}^\nu) = \sum_{k=1}^d \pi_t^\nu(k) \rho_{jk}(t)$, and (2.4) provide

$$\begin{aligned}
|\pi_t^\nu(i) - \pi_t^{\beta\nu}(i)| &\leq \max_{j \in \mathbb{S}} \frac{d\beta}{d\nu}(j) \max_{j \in \mathbb{S}} \frac{d\nu}{d\beta}(j) \sum_{j=1}^d \left| \sum_{k=1}^d \pi_t^\nu(k) \rho_{jk}(t) - \rho_{ji}(t) \right| \\
&= \max_{j \in \mathbb{S}} \frac{d\beta}{d\nu}(j) \max_{j \in \mathbb{S}} \frac{d\nu}{d\beta}(j) \sum_{j=1}^d \left| \sum_{k=1}^d \pi_t^\nu(k) (\rho_{jk}(t) - \rho_{ji}(t)) \right|
\end{aligned}$$

and the result. $\square$

Lemma 2.2. (Lemma 9.5, [4].) *Assume ν is positive. Then, for any $j \in \mathbb{S}$,*

$$\begin{aligned}
\rho_{ji}(0) &= \begin{cases} 1, & j = i \\ 0, & j \neq i \end{cases} \\
\frac{d\rho_{ji}(t)}{dt} &= \sum_{r \neq i} \frac{\lambda_{ri} \pi_t^\nu(r)}{\pi_t^\nu(i)} (\rho_{jr}(t) - \rho_{ji}(t)), \quad i = 1, \dots, d. \quad (2.5)
\end{aligned}$$

The following lemma plays a key role in proving (1.4).

Lemma 2.3. *Assume ν is positive and $\lambda_{ij} > 0$ for all $i \neq j$. Then for any i, j, k , P -a.s.*

$$|\rho_{ji}(t) - \rho_{jk}(t)| \leq \exp \left(-2 \min_{p \neq q} \sqrt{\lambda_{pq} \lambda_{qp}} t \right). \quad (2.6)$$

Proof. Set $\rho_\diamond(t) = \min_{i \in \mathbb{S}} \rho_{ji}(t)$ and $\rho^\diamond(t) = \max_{i \in \mathbb{S}} \rho_{ji}(t)$. Obviously, $\rho_\diamond(0) = 0$ and $\rho^\diamond(0) = 1$. Since processes $\rho_{ji}(t)$, $i = 1, \dots, d$ have continuously differentiable paths, piece-wise constant functions², $i^\diamond(t) = \operatorname{argmax}_{i \in \mathbb{S}} \varrho_{ji}(t)$ and $i_\diamond(t) = \operatorname{argmin}_{i \in \mathbb{S}} \varrho_{ji}(t)$ have a finite number of jumps on any finite time interval. Therefore, paths of $\rho^\diamond(t)$ and $\rho_\diamond(t)$ are absolutely continuous functions the derivatives of which are defined, in accordance to (2.5), as

$$\begin{aligned} \frac{d\rho_\diamond(t)}{dt} &= \sum_{r \neq i_\diamond(t)} \frac{\lambda_{ri_\diamond(t)} \pi_t^\nu(r)}{\pi_t^\nu(i_\diamond(t))} (\rho_{jr}(t) - \rho_\diamond(t)) \\ \frac{d\rho^\diamond(t)}{dt} &= \sum_{r \neq i^\diamond(t)} \frac{\lambda_{ri^\diamond(t)} \pi_t^\nu(r)}{\pi_t^\nu(i^\diamond(t))} (\rho_{jr}(t) - \rho^\diamond(t)). \end{aligned} \quad (2.7)$$

Notice that for any r we have $\rho_{jr}(t) - \rho_\diamond(t) \geq 0$, and $\rho_{jr}(t) - \rho^\diamond(t) \leq 0$, so that by (2.7)

$$\begin{aligned} \frac{d\rho_\diamond(t)}{dt} &\geq \frac{\lambda_{ri_\diamond(t)} \pi_t^\nu(r)}{\pi_t^\nu(i_\diamond(t))} (\rho_{jr}(t) - \rho_\diamond(t)), \quad r \neq i_\diamond(t) \\ \frac{d\rho^\diamond(t)}{dt} &\leq \frac{\lambda_{ri^\diamond(t)} \pi_t^\nu(r)}{\pi_t^\nu(i^\diamond(t))} (\rho_{jr}(t) - \rho^\diamond(t)), \quad r \neq i^\diamond(t). \end{aligned}$$

Consequently, taking $r = i^\diamond(t)$ in the first case and $r = i_\diamond(t)$ for the second one, we get under $i^\diamond(t) \neq i_\diamond(t)$.

$$\begin{aligned} \frac{d\rho_\diamond(t)}{dt} &\geq \frac{\lambda_{i^\diamond(t)i_\diamond(t)} \pi_t^\nu(i^\diamond(t))}{\pi_t^\nu(i_\diamond(t))} (\rho^\diamond(t) - \rho_\diamond(t)) \\ \frac{d\rho^\diamond(t)}{dt} &\leq -\frac{\lambda_{i_\diamond(t)i^\diamond(t)} \pi_t^\nu(i_\diamond(t))}{\pi_t^\nu(i^\diamond(t))} (\rho^\diamond(t) - \rho_\diamond(t)). \end{aligned} \quad (2.8)$$

If $i^\diamond(t) = i_\diamond(t)$, then $\rho^\diamond(t) = \rho_\diamond(t)$ and by (2.8) $\frac{d\rho_\diamond(t)}{dt} = 0$. Hence, (2.8) is always valid.

Set $\Delta_t = \rho^\diamond(t) - \rho_\diamond(t)$ and notice $\Delta_0 = 1$. Owing to (2.8), a differential inequality holds true

$$\frac{d\Delta_t}{dt} \leq -\left(\frac{\lambda_{i^\diamond(t)i_\diamond(t)} \pi_t^\nu(i^\diamond(t))}{\pi_t^\nu(i_\diamond(t))} + \frac{\lambda_{i_\diamond(t)i^\diamond(t)} \pi_t^\nu(i_\diamond(t))}{\pi_t^\nu(i^\diamond(t))} \right) \Delta_t. \quad (2.9)$$

Now, an obvious inequality

$$\frac{\lambda_{i^\diamond(t)i_\diamond(t)} \pi_t^\nu(i^\diamond(t))}{\pi_t^\nu(i_\diamond(t))} + \frac{\lambda_{i_\diamond(t)i^\diamond(t)} \pi_t^\nu(i_\diamond(t))}{\pi_t^\nu(i^\diamond(t))} \geq \min_{x \geq 0} \left(\lambda_{i^\diamond(t)i_\diamond(t)} x + \lambda_{i_\diamond(t)i^\diamond(t)} \frac{1}{x} \right) \quad (2.10)$$

and

$$\min_{x \geq 0} \left(\lambda_{i^\diamond(t)i_\diamond(t)} x + \lambda_{i_\diamond(t)i^\diamond(t)} \frac{1}{x} \right) = 2\sqrt{\lambda_{i^\diamond(t)i_\diamond(t)} \lambda_{i_\diamond(t)i^\diamond(t)}} \geq 2 \min_{p \neq q} \sqrt{\lambda_{pq} \lambda_{qp}}$$

²if max and min are attained on several numbers, the smallest is taken

provide

$$\frac{d\Delta_t}{dt} \leq -2 \min_{p \neq q} \sqrt{\lambda_{pq} \lambda_{qp}} \Delta_t$$

and the result. $\square$

2.2. The proof of (1.4). Since $\|\pi_t^\nu - \pi_t^{\beta\nu}\| = \sum_{i=1}^d |\pi_t^\nu(i) - \pi_t^{\beta\nu}(i)|$, by Lemma 2.1 we have

$$\|\pi_t^\nu - \pi_t^{\beta\nu}\| \leq d^2 \max_{j \in \mathbb{S}} \frac{d\beta}{d\nu} \max_{j \in \mathbb{S}} \frac{d\nu}{d\beta} \max_{1 \leq i, j, k \leq d} |\rho_{ji}(t) - \rho_{jk}(t)|, \quad P - \text{a.s.}$$

and then by Lemma 2.3 the result. $\square$

3. THE PROOF OF (1.3)

1. Assume ν is positive and $\beta' \sim \beta'' \sim \nu$. Let β denote any of β', β'' . Then, due to Theorem 1.1, we have

$$\overline{\lim}_{t \rightarrow \infty} \frac{1}{t} \log \|\pi_t^\nu - \pi_t^{\beta\nu}\| \leq -2 \min_{p \neq q} \sqrt{\lambda_{pq} \lambda_{qp}}, \quad P - \text{a.s.} \quad (3.1)$$

Hence, by the triangular inequality

$$\begin{aligned} \|\pi_t^{\beta'\nu} - \pi_t^{\beta''\nu}\| &\leq \|\pi_t^{\beta'\nu} - \pi_t^\nu\| + \|\pi_t^{\beta''\nu} - \pi_t^\nu\| \\ &\leq 2 \max \left(\|\pi_t^{\beta'\nu} - \pi_t^\nu\|, \|\pi_t^{\beta''\nu} - \pi_t^\nu\| \right) \end{aligned}$$

(1.3) holds true.

2. It can be readily checked that for any fixed $\delta > 0$ the values π_δ^ν and $\pi_\delta^{\beta\nu}$, defined by (1.1) and (2.1) respectively, have positive entries Q^ν -a.s. Now, the original filtering problem, under obvious redefinitions, can be translated into filtering problem on $[\delta, \infty)$ for which (1.3) follows.

REFERENCES

- [1] R. Atar, O. Zeitouni, Lyapunov exponents for finite state nonlinear filtering, *SIAM J. Control Optim.*, Vol. 35, No 1., pp. 36-55, 1997
- [2] Delyon, B., Zeitouni, O. Lyapunov exponents for filtering problems in *Applied Stochastic Analysis*, ed. M. Davis, R. Elliot, pp. 511-521, Gordon and Breach, New York, 1991
- [3] Le Gland, F., Mevel, L., Exponential forgetting and geometric ergodicity in Hidden Markov Models, *Math. Control Signals Systems*, Vol. 13, **1**, pp. 63-93, 2000
- [4] Liptser, R., Shiryaev, A., *Statistics of Random Processes: General Theory*, Springer, 2001
- [5] Wonham, W.M. Some applications of stochastic differential equations to optimal nonlinear filtering. *SIAM J. Control Optimization*, **2**, 347-69, 1965.

DEPT. ELECTRICAL ENGINEERING-SYSTEMS, TEL AVIV UNIVERSITY, 69978 TEL AVIV, ISRAEL

E-mail address: pavelm@eng.tau.ac.il

DEPARTMENT OF ELECTRICAL ENGINEERING-SYSTEMS, TEL AVIV UNIVERSITY, 69978 TEL AVIV, ISRAEL

E-mail address: liptser@eng.tau.ac.il